



If you would like to refer a patient for an Axumin PET/CT scan, please complete this form and fax it to an Axumin imaging site. To locate an Axumin imaging site, visit [www.axumin.com/locator](http://www.axumin.com/locator).

**PATIENT INFORMATION**

Patient Name	Date of Birth	Phone
Primary Insurance	Subscriber ID	Prior Authorization #
Secondary Insurance	Subscriber ID	Prior Authorization #
Referring Physician	Phone	Physician Fax

**CLINICAL SIGNS/SYMPTOMS**

Diagnosis	Clinical Question	ICD-10 Code(s)
-----------	-------------------	----------------

**SPECIFIC REASON FOR AXUMIN PET STUDY**

Detecting suspected recurrence or metastasis of previously treated prostate cancer

Suspected recurrence based on:      Document PSA nadir and subsequent rise:

Special instructions:

**PRIOR STUDIES/TREATMENT**

Radical prostatectomy <input type="checkbox"/> YES <input type="checkbox"/> NO	Physician	Date
Radiation therapy to prostate <input type="checkbox"/> YES <input type="checkbox"/> NO	Physician	Date
Other treatments (describe)	Physician	Date

**PREVIOUS SCANS**

<input type="checkbox"/> CT <input type="checkbox"/> MR <input type="checkbox"/> Bone scan	Location	Date
--	----------	------

I verify that the above information is complete and accurate to the best of my knowledge and that I have prescribed Axumin based on my independent professional judgment of medical necessity. I represent and warrant that I have my patient's authorization on file to (i) disclose his/her health information and to transfer such information to Blue Earth Diagnostics and its agents to use and disclose as necessary to provide reimbursement services.

Physician or Nurse Practitioner Signature	Date
---	------

# AXUMIN®

## Fluciclovine F 18 Injection

### PATIENT PREPARATION AND PRECAUTIONS

#### PREPARING FOR AN AXUMIN SCAN:

- Patient should avoid any significant exercise for at least one day prior to PET/CT imaging
- Patient should fast for at least four (4) hours prior to administration (other than small amounts of water for taking medications)
- The total scan time is between 20 and 30 minutes

#### INDICATION

Axumin® (fluciclovine F 18) injection is indicated for positron emission tomography (PET) imaging in men with suspected prostate cancer recurrence based on elevated blood prostate specific antigen (PSA) levels following prior treatment.

#### IMPORTANT SAFETY INFORMATION

- Image interpretation errors can occur with Axumin PET imaging. A negative image does not rule out recurrent prostate cancer and a positive image does not confirm its presence. The performance of Axumin seems to be affected by PSA levels. Axumin uptake may occur with other cancers and benign prostatic hypertrophy in primary prostate cancer. Clinical correlation, which may include histopathological evaluation, is recommended.
- Hypersensitivity reactions, including anaphylaxis, may occur in patients who receive Axumin. Emergency resuscitation equipment and personnel should be immediately available.
- Axumin use contributes to a patient's overall long-term cumulative radiation exposure, which is associated with an increased risk of cancer. Safe handling practices should be used to minimize radiation exposure to the patient and health care providers.
- Adverse reactions were reported in  $\leq 1\%$  of subjects during clinical studies with Axumin. The most common adverse reactions were injection site pain, injection site erythema and dysgeusia.

To report suspected adverse reactions to Axumin, call 1-855-AXUMIN1 (1-855-298-6461) or contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**Full Axumin prescribing information is available at [www.axumin.com](http://www.axumin.com).**